


Hill, J. M. and V. W. Woodward, Assays for aspartate and ornithine transcarbamylase by means of the pH-stat.

Aspartate transcarbamylase (ATCase) activity has been determined by a calorimetric assay for carbamyl aspartate (Davis and Woodward 1962 Genetics 47: 1075). The colorimetric assay

can be adapted to ornithine transcarbamylase activity by assaying for citrulline. These same enzymes can be assayed by titration of hydrogen ion released in a pH-stat. The following equation describes the reaction in question:



carbamyl phosphate

aspartate

carbamyl aspartate

Reagents: 0.005 M sulfamic acid (primary standard solution); 0.01 M NaOH, standardized against sulfamic acid; 0.05 M L-aspartate, standard solution (0.665 g dissolved in 50 ml 0.10 M NaOH and diluted to 100 ml); 0.005 M carbamyl phosphate (0.0204 g dissolved in 25 ml cold water and kept in ice bath); H₂O, boiled and copped to exclude CO₂; protein solution adjusted to pH 8.5 with NaOH.

Apparatus: Sargent pH-stat equipped as follows: 2.5 ml barrel and plunger; 10 ml reaction container, with stopper permitting entrance of electrodes, thermocouple, thermometer, NaOH delivery tube, nitrogen delivery tube, and reaction delivery syringe. Nitrogen cylinder and passage for delivery of nitrogen through 0.5 M NaOH.

Procedure: The pH-stat is calibrated against standard pH 8.0 buffer at 30°C. The barrel is filled with 0.01 M NaOH and the system is flushed with nitrogen. Water, enzyme and aspartate are introduced into the reaction vessel and the mixture is titrated to pH 8.5. The reaction is begun with the addition of carbomyl phosphate. At the end of the assay an aliquot of 0.005 M sulfamic acid is titrated under the conditions of the assay to standardize the NaOH. From this titration the number of μmoles of H⁺ released during the assay can be calculated. Also, carbamyl phosphate can be assayed by letting the reaction go to completion, i.e., by permitting all the carbomyl phosphate to convert to a carbamyl amino acid, and determining the number of μmoles H⁺ released.

Starting with 2.00 ml 0.05 M aspartate, 0.180 units of enzyme, H₂O, 0.01 M NaOH, 0.50 ml 0.00425 M carbamyl phosphate in a total final volume of 8.5 ml, it was determined that the NaOH was 0.00947 M, and that the initial velocity of the reaction was 0.1843 μmole H⁺ per minute. Subtracting carbomyl phosphate hydrolysis, 0.0057 μmoles H⁺ per minute, leaves an initial velocity of 0.1786 μmoles H⁺ per minute. Aliquots of this reaction mix were assayed colorimetrically, and the initial velocity was shown to be 0.180 μmoles carbamyl aspartate per minute. This close agreement speaks for the validity of the pH-stat assay method at this pH and under these conditions.

This work was performed under USPHS Grant GM-10206 -03. ■ ■ ■ Biology Department, Rice University, Houston, Texas.