

A precise size-estimate for the small RNA products arising from *Neurospora crassa* Dicer activity.

Paul ReFalo and Matthew S. Sachs, Department of Environmental and Biomolecular Systems, Oregon Health & Science University, Beaverton OR 97006-8921

Fungal Genetics Newsletter 51:21-22

Neurospora crassa cell-free extracts prepared from strains containing one or both functional Dicer genes, but not from a strain lacking functional Dicer genes, converts radiolabeled double-strand RNA (dsRNA) in an energy-dependent manner into short RNAs with an estimated size of ~25-nt (Catalanotto *et al.* 2004). A smaller nucleolytic digestion product was also produced in an energy-dependent manner from either dsRNA or single-stranded RNA. Here we obtained more precise sizes for these products by electrophoresis of samples on a long (40-cm) denaturing DNA sequencing gel (20% polyacrylamide/7M urea).




Fig. 1. Dicer activity in *N.crassa* single and double mutants for *dcl-1* and *dcl-2*. Analysis of WT, *dcl-1*, *dcl-2* and *dcl-1/dcl-2* strains for Dicer activity was accomplished as described (Catalanotto *et al.* 2004). *N. crassa* cell-free extracts were incubated with radiolabeled dsRNA for 0, 30, or 90 minutes (T_0 , T_{30} , T_{90}) in the presence or absence of an energy regenerating system as indicated, and the RNA was examined by denaturing gel electrophoresis on a denaturing 20% polyacrylamide sequencing gel. Decade RNA Markers (Ambion) labeled with ^{32}P were used as size standards (lanes marked M).

The data (Fig. 1) show the Dicer- and energy-dependent products obtained from the dsRNA substrate were clustered in a region indicating the majority of species had sizes between 21-26 nt, with most approximately 23-nt in length. This is

consistent with results from other organisms (Agrawal *et al.* 2003). An additional very small degradation product was produced in an energy-dependent but Dicer-independent manner. This product, but not the ~21-26 nt products, was also obtained from single strand RNA (data not shown). Apparently, this small product migrated anomalously (suggesting a size of approximately ~16-nt) in the shorter gels containing a lower-percentage of polyacrylamide that were used previously (Catalanotto *et al.* 2004). Thus, the results in Fig. 1 indicate that, from input dsRNA, *N. crassa* extracts with Dicer activity produced RNAs of the size expected to function as small interfering RNA. *N. crassa* extracts did not contain other activities that processed input RNA into other large oligonucleotide products.

Acknowledgment. This work was supported by the NIH (GM47498).

References

- Agrawal, N., P. V. Dasaradhi, A. Mohammed, P. Malhotra, R. K. Bhatnagar and S. K. Mukherjee, 2003. RNA interference: biology, mechanism, and applications. *Microbiol. Mol. Biol. Rev.* 67: 657-685.
- Catalanotto, C., M. Pallotta, P. Refalo, M. S. Sachs, L. Vayssie, G. Macino and C. Cogoni, 2004. Redundancy of the two dicer genes in transgene-induced posttranscriptional gene silencing in *Neurospora crassa*. *Mol. Cell. Biol.* 24: 2536-2545.