

Neurospora genetics at the turn of the century

David D. Perkins, Department of Biological Sciences, Stanford University, Stanford, California 94305-5020

Research advances are described that have been made since a 1992 survey in *Genetics* (130: 687-701).

The century's end also marks the 75th anniversary of the first genetic experiments with *Neurospora*. Although accounts usually begin with the 1941 paper of Beadle and Tatum, genetic analysis was in fact initiated 16 years earlier by B. O. Dodge. It was he who identified the two mating types and used them to demonstrate mendelian segregation in individual asci. Dodge quickly recognized the potentialities of the organism. His enthusiasm was largely responsible for the adoption of *Neurospora* by geneticists and for its development as a model organism. Now, 75 years after Dodge's identification of the first gene, over 1000 loci have been characterized and mapped in the seven linkage groups. Horowitz (1991), Perkins (1992), and Davis (2000) have described the main long-term contributions of *Neurospora* and have evaluated their significance. Specific advances since the 1992 review are summarized below.

Neurospora crassa has become a preeminent model species for studying circadian rhythms. In an extension of work pioneered by J. F. Feldman, the gene *frq* (frequency) has been shown to encode a central component of a molecular feedback loop in which the product of *frq* negatively regulates synthesis of its own transcript, resulting in daily oscillation (Aronson *et al.* 1994, Dunlap 1993). A variety of clock-controlled genes have been identified, the inactivation of which does not alter rhythmicity (Bell-Pedersen *et al.* 1996). Resetting the clock occurs when induction of *frq* by light overcomes negative autoregulation, resulting in phase delay or advance, depending on the time of day (Crosthwaite *et al.* 1995). Photoreceptor regulator genes (*white-collar-1* and *-2*) are essential in assembly or operation of the *frq* feedback loop (Crosthwaite *et al.* 1997). Demonstration that entrainable and free running rhythmicity can persist in the absence of *frq* and *wc* gene-products has suggested increasingly sophisticated models (Morrow *et al.* 1999, Lakin-Thomas and Brody 2000, McWatters *et al.* 1999). For reviews see Bell-Pedersen (1998), Loros (1998), Dunlap (1999), Lakin-Thomas (2000).

Extensive information has been obtained on genes the expression of which is under photo-, circadian, or developmental control. (See Lauter 1996, Bell-Pedersen *et al.* 1996, Ebbole 1995). Genes have been isolated and characterized that specify α -subunits of heterotrimeric GTP binding proteins (Turner and Borkovich 1993, Baasiri *et al.* 1997, Kays *et al.* 1998). Numerous genes that encode putative signal transduction proteins have been identified (Margolis and Yanofsky 1998).

Significant contributions have been made to the molecular genetics of fungal photobiology, with the identification and characterization of photomutants and genes regulated by blue light (reviewed by Lauter 1996). The two *white collar* genes are global regulators of photoresponses, encoding blue-light-activated transcription factors and participating in the blue-light signal transduction pathway (Ballario and Macino 1997, Schwerdtfeger and Linden 2000). A gene homologous to archaeal rhodopsins provides the first example of an opsin in eukaryotes other than animals; the gene-product is a photochemically reactive member of the archaeal rhodopsin family (Bieszke *et al.* 1999a).

The ascus-dominant expression of the ascospore-maturation gene *Asm-1* has been shown to result from failure of transvection, wherein chromosome rearrangements or ectopic placement of a gene disrupts pairing of allelic chromosomal genes during the sexual phase and results in a mutant phenotype, even in spores that carry the normal allele (Aramayo and Metzberg 1996). This discovery provided a clear demonstration that transvection occurs in an organism very different from *Drosophila*.

Study of the UV-sensitive mutant *mus-18* has identified a novel DNA endonuclease that initiates an excision repair pathway completely different from previously known DNA-repair mechanisms (Ishii *et al.* 1991; see Yasui and McCready 1998 for review). A UV-sensitive mutant, *mus-38*, is impaired in the previously known highly conserved nucleotide excision repair pathway (Ishii *et al.* 1998).

Evidence has accumulated that RIP (repeat-induced point mutation) serves as a genome defense system (see Selker 1997). While only one active transposon has been found in *Neurospora*, sequences have been discovered that represent several different transposon families, with unmistakable hallmarks of RIP (Cambareri *et al.* 1998, Kinsey *et al.* 1994, Margolin *et al.* 1998, Bibbins and Connerton 1998). Characterization of centromeric DNA (Centola and Carbon 1994) has revealed the presence of complex repeats reminiscent of the centric heterochromatin of *Drosophila* (Cambareri *et al.* 1998). Defective transposable elements of several types are present among the repeats, and these show evidence of having been inactivated by RIP.

RIP has been used extensively for gene disruption. Null mutations of RIP-inactivated essential genes can be recovered by using a meiotic mutant that produces heterokaryotic ascospores (Metzberg and Grotelueschen 1992; Harkness *et al.* 1994).

RIP was shown frequently to generate signals for de novo methylation. Evidence was also obtained for maintenance methylation in *Neurospora* (Singer *et al.* 1995). Further analysis of methylation resulting from RIP led to the discovery of an unexpected connection between protein acetylation and DNA methylation (Selker 1998). Mutants defective in DNA methylation (*dim* mutants) have been isolated (see Foss *et al.* 1998). Mutations in *dim-2*, which is thought to encode a DNA methyltransferase (E. Kouzminova and E. U. Selker, personal communication), result in loss of all detectable methylation, at least in the vegetative phase. (No known mutation in any other eukaryote completely abolishes DNA methylation.) Identification of *dim-2* indicated that DNA methylation is not essential in *Neurospora*. The mutant has been used to demonstrate that methylation can either interfere

with gene expression (Irelan and Selker 1997, Rountree and Selker 1997) or promote it indirectly (Cambareri *et al.* 1996), that methylation can inhibit transcript elongation *in vivo* (Rountree and Selker 1997), and that gene silencing in the vegetative phase ("quelling") does not rely on DNA methylation (Cogoni *et al.* 1996).

Reversible silencing (quelling) can occur when additional copies of a gene are introduced by transformation (Rom and Macino 1992, Pandit and Russo 1992; reviewed by Irelan and Selker 1996). Both the introduced and the resident copies are affected. Silencing is posttranscriptional and is dominant in heterokaryons (see Cogoni *et al.* 1996, Cogoni and Macino 1997, and references therein). Quelling-deficient mutants in which transgene-induced gene silencing is impaired have been used to show that quelling requires a protein homologous to RNA-dependent RNA polymerase (Cogoni and Macino 1999a, a RecQ DNA helicase (known to be involved in repair and recombination in other organisms) (Cogoni and Macino 1999b), and a homolog of *C. elegans rde-1*, which controls the degradation of double-stranded RNA (Catalanotto *et al.* 2000).

Mitochondrial tRNA synthetase has been shown to mediate RNA self-splicing. Two mitochondrial plasmids are retroelements that share properties of RNA viruses and mitochondrial introns. The novel transcriptases they encode possess characteristics suggesting how present-day reverse transcriptases and DNA polymerases could have evolved (Wang and Lambowitz 1993).

Regulated ribosome stalling has been demonstrated (Wang and Sachs 1997).

Vesicles from the outer mitochondrial membrane have been purified on a massive scale and the preprotein translocase (TOM complex) has been shown by electron microscopy to contain centers interpreted as pores that represent protein-conducting channels (Künkele *et al.* 1998).

Over 4600 cultures from natural populations throughout the world are now available for study (Turner and Perkins 2000). Wild-collected strains have provided information on species distribution, ecology, genetic diversity, population structure, and meiotic drive systems. They have also been a source of genetic variants for a variety of laboratory investigations.

Surveys of strains from nature have revealed the widespread occurrence of mitochondrial plasmids, which belong to discrete families (Yang and Griffiths 1993, Arganoza *et al.* 1994). New examples have been discovered of plasmids that cause senescence (Yang and Griffiths 1993, Marcinko-Kuehn *et al.* 1994, He *et al.* 2000; reviewed by Griffiths 1992, 1993, 1998).

Investigations with the pseudohomothallic species *N. tetrasperma* have revealed novel features of this unique genetic system (Merino *et al.* 1996, Gallegos *et al.* 2000, Metzenberg and Randall 1995, Raju and Perkins 1994).

Integration of transforming DNA was shown to be accompanied by new gross chromosome rearrangements, many of which have breakpoints associated with vector DNA (Perkins *et al.* 1993).

Morphological mutants called *ropy* were shown to be defective in specifying subunits of dynein and related molecular motors (Plamann *et al.* 1994). Mutations at *ropy* loci are selectable as suppressors of the morphological mutant *cot-1*. Similarly, *mcb* (*microcycle blastoconidiation*) acts as a suppressor of the morphological mutant *crisp* (Bruno *et al.* 1996). Mutations of *mcb* affect growth polarity. Secretion of extracellular enzymes in *mcb* cultures is increased to the high level that is characteristic of the hyphal tip in wild type cultures (Lee *et al.* 1998).

Understanding of meiotic recombination has been advanced by high-resolution experiments using molecular markers (e.g., T. Randall and D. R. Stadler, in preparation). The recombinator gene *cog* has been cloned and two alleles have been sequenced (Yeadon and Catchside 1995). Intragenic recombination appears to be initiated at *cog*⁺ (Yeadon and Catchside 1998), which is 3' of the *am* locus (Bowring and Catchside 1991). Intragenic recombination has been studied using simultaneously both closely linked RFLP markers and more distant classical genes to flank the *am* gene (Bowring and Catchside 1996, 1998) and the *his-3* gene (Yeadon and Catchside 1998). In both cases, conversion tracts frequently are interrupted. Although about one third of gene conversions at *his-3* are accompanied by a crossover, this apparent association is tenuous at *am* where recombination frequencies are much lower. This observation casts doubt on the widely held assumption that both conversion and reciprocal crossing over arise from the same event. Evidence has been obtained that conversion events at *am* stimulate crossing over nearby (Bowring and Catchside 1999). Studies with closely linked molecular markers show that the genetic criteria previously used to establish the order of intragenic sites is flawed when differentially spaced conventional mutants are used as flanking markers (Bowring and Catchside 1995).

Substantial progress has been made in understanding the organization and function of genes at the *mating type* locus (now called *idiomorphs* in recognition of their lack of homology). The *mat a* idiomorph contains a single open reading frame, while *mat A* contains three (Ferreira *et al.* 1996). For reviews see Staben (1996), Coppin *et al.* (1997). Both *mat A-1* and *mat a-1* appear to be essential for mating and for sexual development, while *mat A-2* and *mat A-3* increase fecundity but are not essential (Ferreira *et al.* 1998).

Genes responsible for vegetative (heterokaryon) incompatibility (*het* genes) have been cloned and sequenced (Saupe *et al.* 1996, Smith *et al.* 1996, 1999, 2000; Shiu and Glass 1999). The same multiple alleles of *het-c* that are found in *N. crassa* are also present in other *Neurospora* species and in related genera, indicating that they were derived from a common ancestor and have been conserved during evolution (Wu *et al.* 1998).

The *tol* gene has been cloned and sequenced (Shiu *et al.* 1999). A functional allele of *tol* (*tolerant*) is required for expression of the *mating-type* mediated vegetative incompatibility phenotype that results when *mat A* and *mat a* idiomorphs are together in heterokaryons or heterozygous partial diploids. Vegetative incompatibility reactions mediated by genes other than

mating type do not require the presence of a functional *tol* allele (Leslie and Yamashiro 1998). An active *tol* allele is normally present in the heterothallic outbreeding species *N. crassa*, and the gene was originally identified in that species as a recessive mutant that suppresses *A + a* vegetative incompatibility. The species *N. tetrasperma*, which normally exists as a self-fertile (*mat A + mat a*) heterokaryon, was shown to possess an inactive *tol* allele (Jacobson 1992). The active and inactive *tol* alleles have been interchanged between *N. crassa* and *N. tetrasperma*.

Genes that served as morphological markers in constructing the first fungal genetic maps in the 1930's have now been cloned, sequenced, and characterized functionally (e.g., *crisp-1* (Kore-Eda *et al.* 1991), *fluffy* (Bailey and Ebbole 1998).

Heterokaryons are being used to produce heterodimeric molecules that incorporate components originating from genetically different nuclei. Intact antibody molecules are formed by heterokaryons in which the light chain is produced and secreted by one nuclear type and the heavy chain is produced and secreted by the other (Stuart 1997, 1998).

Genetic mapping has progressed substantially, using both classical and RFLP markers (Perkins 2000, Nelson *et al.* 1998, Nelson and Perkins 2000).

Genome projects are under way. Expressed sequence tags (ESTs) have been obtained that identify genes expressed at different stages of the life cycle or during different parts of the circadian cycle. More than half the expressed sequences show no similarity to genes previously identified in the yeast genome or elsewhere. Over 2000 different genes have been identified in this way (Nelson *et al.* 1997, Dolan *et al.* 2000). Over 50% of these have no known homologs in any organism (Nelson and Narvig 1999). Physical maps of the genome are being constructed (Arnold, 2000), and DNA sequencing of the genome is progressing (Mewes *et al.* 2000; <http://www.mips.biochem.mpg.de/proj/neurospora/>).

At the turn of the century, *Neurospora* is genetically and biologically the best known eukaryotic microorganism. The rich resources of information, brought together by Davis (2000) and by Perkins *et al.* (2000), will speed progress in relating sequence data to biologically meaningful problems.

REFERENCES

- Aramayo, R., and R. L. Metzenberg, 1996 Meiotic transvection in fungi. *Cell* 86: 103-113.
- Arganoza, M. T., J. Min, Z. Hu and R. A. Akins, 1994 Distribution of seven homology groups of mitochondrial plasmids in *Neurospora*: Evidence for widespread mobility between species in nature. *Curr. Genet* 26: 62-73.
- Arnold, J., 2000 Physical map of *Neurospora crassa*. *Fungal Genet. Newslett.* 47S: 16. (Abstr.)
- Aronson, B. D., K. A. Johnson, J. J. Loros and J. C. Dunlap, 1994 Negative feedback defining a circadian clock: Autoregulation of the clock gene *frequency*. *Science* 263: 1578-1584.
- Ballario, P., and G. Macino, 1997 White-collar proteins: PASSing the light signal in *Neurospora crassa*. *Trends Microbiol.* 5: 458-462.
- Baasiri, R. A., X. Lu, P. S. Rowley, G. E. Turner and K. A. Borkovich, 1997 Overlapping functions for two G protein α -subunits in *Neurospora crassa*. *Genetics* 147: 137-145.
- Bailey, L. A., and D. J. Ebbole, 1998 The *fluffy* gene of *Neurospora crassa* encodes a Gal4p-type C6 zinc cluster protein required for conidial development. *Genetics* 148: 1813-1820.
- Bell-Pedersen, D., 1996 Keeping pace with *Neurospora* circadian rhythms. *Microbiology* 144: 1699-1711.
- Bell-Pedersen, D., M. Shinohara, J. J. Loros and J. C. Dunlap, 1996 Clock-controlled genes isolated from *Neurospora crassa* are late night- to morning-specific. *Proc. Nat. Acad. Sci. U.S.A.* 93: 13096-13101.
- Bieszke, J. A., E. L. Braun, L. E. Bean, S. Kang, D. O. Narvig, and K. A. Borkovich, 1999a The *nop-1* gene of *Neurospora crassa* encodes a seven-helix transmembrane retinal-binding protein homologous to archaeal rhodopsins. *Proc. Nat. Acad. Sci. U.S.A.* 96: 8034-8039.
- Bieszke, J. A., E. N. Spudich, K. L. Scott, K. A. Borkovich and J. L. Spudich, 1999b A eukaryotic protein, NOP-1, binds retinal to form an archaeal rhodopsin-like photochemically reactive pigment. *Biochemistry* 38: 14138-14145.
- Bowring, F. J., and D. E. A. Catcheside, 1991 The initiation site for recombination *cog* is at the 3' end of the *his-3* gene in *Neurospora crassa*. *Mol. Gen. Genet.* 229: 273-277.
- Bowring, F. J., and D. E. A. Catcheside, 1995 The orientation of gene maps by recombination of flanking markers for the *am* locus of *Neurospora crassa*. *Curr. Genet.* 29: 27-33.
- Bowring, F. J., and D. E. A. Catcheside, 1996 Gene conversion alone accounts for more than 90% of recombination events at the *am* locus of *Neurospora crassa*. *Genetics* 143: 129-136.
- Bowring, F. J., and D. E. A. Catcheside, 1998 Analysis of conversion tracts associated with recombination events at the *am* locus in *Neurospora crassa*. *Curr. Genet.* 29: 43-49.
- Bowring, F. J., and D. E. A. Catcheside, 1999 Evidence for negative interference: Clustering of crossovers close to *am* amongst *am* recombinants. *Genetics* 152: 965-969.
- Bruno, K. S., R. Aramayo, P. F. Minke, R. L. Metzenberg and M. Plamann, 1996 Loss of growth polarity and mislocalization of septa in a *Neurospora* mutant altered in the regulatory subunit of cAMP-dependent protein kinase. *EMBO J.* 15: 5772-5782.

- Cambareri, E. B., R. Aisner and J. Carbon, 1998 Structure of the chromosome VII centromere region in *Neurospora crassa*: Degenerate transposons and simple repeats. *Mol. Cell Biol.* 18: 5465-5477.
- Cambareri, E. B., H. M. Foss, M. R. Rountree, E. U. Selker and J. A. Kinsey, 1996 Epigenetic control of a transposon-inactivated gene in *Neurospora* is dependent on DNA methylation. *Genetics* 143: 137-146.
- Catalanotto, C., G. Azzalin, G. Macino and C. Cogoni, 2000 Transcription: Gene silencing in worms and fungi. *Nature* 404: 245.
- Centola, M., and J. Carbon, 1994 Cloning and characterization of centromeric DNA from *Neurospora crassa*. *Mol. Cell Biol.* 14: 1510-1519.
- Cogoni, C., and G. Macino, 1997 Isolation of quelling-defective (*qde*) mutants impaired in posttranscriptional transgene-induced gene silencing in *Neurospora crassa*. *Proc. Nat. Acad. Sci. U.S.A.* 94: 10233-10238.
- Cogoni, C., and G. Macino, 1999a Gene silencing in *Neurospora crassa* requires a protein homologous to RNA-dependent RNA polymerase. *Nature* 399: 166-169.
- Cogoni, C., and G. Macino, 1999b Posttranscriptional gene silencing in *Neurospora* by a *recQ* DNA helicase. *Science* 286: 2342-2344.
- Cogoni, C., J. T. Irelan, M. Schumacher, T. J. Schmidhauser, E. U. Selker and G. Macino, 1996 Transgene silencing of the *al-1* gene in vegetative cells of *Neurospora* is mediated by a cytoplasmic effector and does not depend on DNA-DNA interactions or DNA methylation. *EMBO J.* 15: 3153-3163.
- Coppin, E., R. Debuchy, S. Arnaise and M. Picard, 1997 Mating types and sexual development in filamentous ascomycetes. *Microbiol. Mol. Biol. Rev.* 61: 411-428.
- Crosthwaite, S. K., J. J. Loros and J. C. Dunlap, 1995 Light-induced resetting of a circadian clock is mediated by a rapid increase in *frequency* transcript. *Cell* 81: 1003-1012.
- Crosthwaite, S. K., J. J. Loros and J. C. Dunlap, 1997 *Neurospora wc-1* and *wc-2*: Transcription, photoresponses, and the origins of circadian rhythmicity. *Science* 226: 763-769.
- Davis, R. W., 2000 *Neurospora: Contributions of a Model Organism*. Oxford University Press.
- Dolan, P. L., D. O. Natvig and M. A. Nelson, 2000 *Neurospora* proteome 2000. *Fungal Genet. Newslett.* 47.
- Dunlap, J. C., 1999 Molecular bases for circadian clocks. *Cell* 96: 271-290.
- Ebbole, D., 1996 Morphogenesis and vegetative differentiation in filamentous fungi. *J. Genet.* 75: 361-374.
- Ferreira, A. V. B., S. Saupé and N. L. Glass, 1996 Transcriptional analysis of the *mt A* idiomorph of *Neurospora crassa* identifies two genes in addition to *mt A-1*. *Mol. Gen. Genet.* 250: 767-774.
- Ferreira, A. V.-B., Z. An, R. L. Metzberg and N. L. Glass, 1998 Characterization of *mat A-2*, *mat A-3*, and Δ *mat A* mating-type mutants of *Neurospora crassa*. *Genetics* 148: 1069-1079.
- Foss, H. M., C. J. Roberts and E. U. Selker, 1998 Reduced levels and altered patterns of DNA methylation caused by mutations in *Neurospora crassa*. *Mol. Gen. Genet.* 259: 60-71.
- Gallegos, A., D. J. Jacobson, N. B. Raju, M. P. Skupski and D. O. Natvig, 2000 Suppressed recombination and a pairing anomaly on the mating-type chromosome of *Neurospora tetrasperma*. *Genetics* 154: 623-633.
- Griffiths, A. J. F., 1992 Fungal senescence. *Annu. Rev. Genet.* 26: 351-372.
- Griffiths, A. J. F., 1995 Natural plasmids of filamentous fungi. *Microbiol. Rev.* 59: 673-685.
- Griffiths, A. J. F., 1998 The kalilo family of fungal plasmids. *Bot. Bull. Acad. Sinica* 39: 147-152.
- Harkness, T. A. A., R. L. Metzberg, H. Schneider, R. Lill, W. Neupert and F. E. Nargang, 1994 Inactivation of the *Neurospora crassa* gene encoding the mitochondrial protein import receptor MOM19 by the technique of "sheltered RIP". *Genetics* 136: 107-118.
- He, C., N. de Groot, J. W. Bok and A. J. F. Griffiths, 2000 Kalilo plasmids are a family of four distinct members with individual global distributions across species. *Curr. Genet.* 37: 39-44.
- Horowitz, N. H., 1991 Fifty years ago: The *Neurospora* revolution. *Genetics* 127: 631-635.
- Irelan, J. T., and E. U. Selker, 1996 Gene silencing in filamentous fungi: RIP, MIP and quelling. *J. Genet.* 75: 313-324.
- Irelan, J. T., and E. U. Selker, 1997 Cytosine methylation associated with repeat-induced point mutation causes epigenetic gene silencing in *Neurospora crassa*. *Genetics* 146: 509-523.
- Ishii, C., K. Nakamura and H. Inoue, 1991 A novel phenotype of an excision-repair mutant in *Neurospora crassa*: Mutagen sensitivity of the *mus-18* mutant is specific to UV. *Mol. Gen. Genet.* 228: 33-39.
- Ishii, C., K. Nakamura and H. Inoue, 1998 A new UV-sensitive mutant that suggests a second excision repair pathway in *Neurospora crassa*. *Mutat. Res.* 408: 171-182.
- Jacobson, D. J., 1992 Control of mating type heterokaryon incompatibility by the *tol* gene in *Neurospora crassa* and *N. tetrasperma*. *Genome* 35: 347-353.
- Kays, A. M., P. S. Rowley, R. Baasiri and K. A. Borkovich, 1998 A G protein subunit, *gna-3*, regulates cAMP metabolism & aerial hyphae formation in *Neurospora crassa*. *Neurospora 1998* (*Neurospora* Information Conference, Asilomar, California), p 22. (Abstr.)

- Kinsey, J. A., P. W. Garrett-Engle, E. B. Cambareri and E. U. Selker, 1994 The *Neurospora* transposon *Tad* is sensitive to repeat-induced point mutation (RIP). *Genetics* 138: 657-664.
- Kore-eda, S., T. Murayama and I. Uno, 1991 Isolation and characterization of the adenylate cyclase structural gene of *Neurospora crassa*. *Jpn. J. Genet.* 66: 317-334.
- Kunkele, K.-P., S. Heins, M. Derabowski, F. E. Nargang, R. Benz, M. Thieffry, J. Walz, R. Lill, S. Nussberger and W. Neupert, 1998 The preprotein translocation channel of the outer membrane of mitochondria. *Cell* 93: 1009-1019.
- Lakin-Thomas, P. L., 2000 Circadian rhythms: New functions for old clock genes? *Trends Genet.* 16: 135-142.
- Lakin-Thomas, P. L., and S. Brody, 2000 Circadian rhythms in *Neurospora crassa*: Lipid deficiencies restore robust rhythmicity to null *frequency* and *white-collar* mutants. *Proc. Nat. Acad. Sci. U.S.A.* 97: 256-261.
- Lauter, F.-R., 1996 Molecular genetics of fungal photobiology. *J. Genet.* 75: 375-386.
- Lee, H., R. G. Walline and M. Plamann, 1998 Apolar growth of *Neurospora crassa* leads to increased secretion of extracellular proteins. *Mol. Gen. Genet.* 29: 209-218.
- Leslie, J. F., and C. T. Yamashiro, 1997 Effects of the *tol* mutation on allelic interactions at *het* loci in *Neurospora crassa*. *Genome* 40: 834-840.
- Loros, J. J., 1998 Time at the end of the millenium: The *Neurospora* clock. *Curr. Opinion Microbiol.* 1: 698-706.
- Marcinko-Kuehn, M., X. Yang, F. Debets, D. J. Jacobson and A. J. F. Griffiths, 1994 A kalilo-like linear plasmid in Louisiana field isolates of the pseudohomothallic fungus *Neurospora tetrasperma*. *Curr. Genet.* 25: 336-343.
- Margolin, B. S., P. W. Garrett-Engle, J. N. Stevens, D. Yen-Fritz, C. Garrett-Engle, R. L. Metzberg and E. U. Selker, 1998 A methylated *Neurospora* 5S rRNA pseudogene contains a transposable element inactivated by RIP. *Genetics* 149: 1787-1797.
- Margolis, P., and C. Yanofsky, 1998 Reverse genetic analysis of signal transduction loci of *N. crassa*. *Neurospora 1998* (*Neurospora* Information Conference, Asilomar, California), p. 24. (Abstr.)
- McWatters, H., J. C. Dunlap and A. J. Millar, 1999 Circadian biology: Clocks for the real world. *Curr. Biol.* 9:R633-R635.
- Merino, S. T., M. A. Nelson, D. J. Jacobson and D. O. Natvig, 1996 Pseudohomothallism and evolution of the mating-type chromosome in *Neurospora tetrasperma*. *Genetics* 143: 789-799.
- Morrow, M., M. Brunner and T. Roenneberg, 1999 Assignment of circadian function for the *Neurospora* clock gene *frequency*. *Nature* 399: 584-586.
- Metzenberg, R. L., and J. S. Grotelueschen, 1992 Disruption of essential genes in *Neurospora* by RIP. *Fungal Genet. Newsl.* 39: 37-49.
- Metzenberg, R. L., and T. A. Randall, 1995 Mating type in *Neurospora* and closely related ascomycetes: Some current problems. *Can. J. Bot.* 73 (Suppl. 1): 5251-5257.
- Mewes H. W., D. Frishman, C. Gruber, B. Geier, D. Haase, A. Kaps, K. Lemcke, G. Mannhaupt, F. Pfeiffer, C. Schuller, S. Stocker and B. Weil, 2000 MIPS: A database for genomes and protein sequences. *Nucl. Acids Res.* 28: 37-40.
- Nelson, M. A., and D. O. Natvig, 2000 The *Neurospora* genome project at UNM: Gold from the mold. *Fungal Genet. Newsl.* 47S:16. (Abstr.)
- Nelson, M. A., and D. D. Perkins, 2000 Restriction polymorphism maps of *Neurospora crassa*: 2000 update. *Fungal Genet. Newsl.* 47.
- Nelson, M. A., S. Kang, E. L. Braun, M. E. Crawford, P. L. Dolan, P. M. Leonard, J. Mitchell, A. M. Armijo, L. Bean, E. Blueyes, T. Cushing, A. Errett, M. Fleharry, M. Gorman, K. Judson, R. Miller, J. Ortega, I. Pavlova, J. Perea, S. Todisco, R. Trujillo, J. Valentine, A. Wells, M. Werner-Washburne, S. Yazzie and D. O. Natvig, 1997 Expressed sequences from conidial, mycelial, and sexual stages of *Neurospora crassa*. *Fungal Genet. Biol.* 21: 348-363.
- Nelson, M. A., M. E. Crawford and D. O. Natvig, 1998 Restriction polymorphism maps of *Neurospora crassa*: 1998 update. *Fungal Genet. Newsl.* 45: 44-54.
- Pandit, N. N., and V. E. A. Russo, 1992 Reversible inactivation of a foreign gene, *hph*, during the asexual cycle in *Neurospora crassa* transformants. *Mol. Gen. Genet.* 234: 412-422.
- Perkins, D. D., 1992 *Neurospora*: The organism behind the molecular revolution. *Genetics* 130: 687-701.
- Perkins, D. D., 2000 *Neurospora crassa* genetic maps and mapped loci. *Fungal Genet. Newsl.* 47.
- Perkins, D. D., J. A. Kinsey, D. K. Asch and G. D. Frederick, 1993 Chromosome rearrangements recovered following transformation of *Neurospora crassa*. *Genetics* 134: 729-736.
- Perkins, D. D., A. Radford and M. S. Sachs, 2000 *The Neurospora Compendium: Chromosomal Loci*. Academic Press.
- Plamann, M., P. F. Minke, J. H. Tinsley and K. S. Bruno, 1994 Cytoplasmic dynein and actin-related protein Arp1 are required for normal nuclear distribution in filamentous fungi. *J. Cell Biol.* 127: 139-149.
- Raju, N. B., and D. D. Perkins, 1994 Diverse programs of ascus development in pseudohomothallic species of *Neurospora*, *Gelasinospora*, and *Podospira*. *Develop. Genet.* 15: 104-118.
- Romano, N., and G. Macino, 1992 Quelling: Transient inactivation of gene expression in *Neurospora crassa* by transformation with homologous sequences. *Mol. Microbiol.* 6: 3343-3353.
- Rountree, M. R., and E. U. Selker, 1997 DNA methylation inhibits elongation but not initiation of transcription in *Neurospora*

crassa. Genes Devel. 11: 2382-2395

- Saupe, S. J., G. A. Kuldau, M. L. Smith and N. L. Glass, 1996 The product of the *het-c* heterokaryon incompatibility gene of *Neurospora crassa* has characteristics of a glycine-rich cell wall protein. Genetics 143: 1589-1600.
- Schwerdtfeger, C., and H. Linden, 2000 Localization and light-dependent phosphorylation of white collar 1 and 2, the two central components of blue light signaling in *Neurospora crassa*. Eur. J. Biochem. 267: 414-421.
- Selker, E. U., 1997 Epigenetic phenomena in filamentous fungi: Useful paradigms or repeat-induced confusion? Trends Genet. 13: 296-301.
- Selker, E. U., 1998 Trichostatin A causes selective loss of DNA methylation in *Neurospora*. Proc. Nat. Acad. Sci. U.S.A. 95: 9430-9435.
- Shiu, P. K. T., and N. L. Glass, 1999 Molecular characterization of *tol*, a mediator of mating-type associated vegetative incompatibility in *Neurospora crassa*. Genetics 151: 545-555.
- Singer, M. J., B. A. Marcotte and E. U. Selker, 1995 DNA methylation associated with repeat-induced point mutation in *Neurospora crassa*. Mol. Cell. Biol. 15: 5586-5597.
- Smith, M. L., C. J. Yang, R. L. Metzenberg and N. L. Glass, 1996 Escape from *het-6* incompatibility in *Neurospora crassa* partial diploids involves preferential deletion within the ectopic segment. Genetics 144: 523-531.
- Smith, M. L., S. P. Hubbard, D. J. Jacobson, O. C. Micali and N. L. Glass, 2000 An osmotic-remedial, temperature sensitive mutation in the allosteric activity site of ribonucleotide reductase in *Neurospora crassa*. Mol. Gen. Genet. 262: 1022-1035.
- Smith, M. L., O. C. Micali, S. P. Hubbard, N. Mir-Rashed, D. J. Jacobson and N. L. Glass, 2000 Vegetative incompatibility in the *het-6* region of *Neurospora crassa* is mediated by two linked genes. Genetics (in press).
- Staben, C., 1996 The mating-type locus of *Neurospora crassa*. J. Genet. 75: 341-350.
- Stuart, W. D., 1997 Heterologous dimeric proteins produced in heterokaryons. U. S. Patent 5,643,745.
- Stuart, W. D., 1998 Novel uses for *Neurospora* heterokaryons. *Neurospora 1998* (Neurospora Information Conference, Asilomar, California). p.9. (Abstr.).
- Turner, B. C., and D. D. Perkins, 1999 *Neurospora* from natural populations: A global study. (In preparation).
- Turner, G. E., and K. A. Borkovich, 1993 Identification of a G protein α -subunit from *Neurospora crassa* that is a member of the G_i family. J. Biol. Chem. 268: 14805-14811.
- Wang, H., and A. M. Lambowitz, 1993 The Mauriceville plasmid reverse transcriptase can initiate cDNA synthesis de novo and may be related to reverse transcriptase and DNA polymerase progenitor. Cell 75: 1071-1081.
- Wang, Z., and M. S. Sachs, 1997 Ribosome stalling is responsible for arginine-specific translational attenuation in *Neurospora crassa*. Mol. Cell. Biol. 17: 4904-4913.
- Wu, J., S. J. Saupe and N. L. Glass, 1998 Evidence for balancing selection operating at the *het-c* heterokaryon incompatibility locus in a group of filamentous fungi. Proc. Nat. Acad. Sci. U.S.A. 95: 12398-12403.
- Yang, X., and A. J. F. Griffiths, 1993 Plasmid diversity in senescent and nonsenescent strains of *Neurospora*. Mol. Gen. Genet. 237: 177-186.
- Yasui, A., and S. J. McCready, 1998 Alternative repair pathways for UV-induced DNA damage. BioEssays 20: 291-297.
- Yeadon, P. J., and D. E. A. Catcheside, 1995 The chromosomal region which includes the recombinator *cog* in *Neurospora crassa* is highly polymorphic. Curr. Genet. 28: 155-163.
- Yeadon, P. J., and D. E. A. Catcheside, 1998 Long interrupted conversion tracts initiated by *cog* in *Neurospora crassa*. Genetics 148: 113-122.