

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 578

1996

Performance of finishing steers offered magnesium-mica in the feedlot ration

K.P. Coffey

F.K. Brazle

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Coffey, K.P. and Brazle, F.K. (1996) "Performance of finishing steers offered magnesium-mica in the feedlot ration," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.1981>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1996 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



**PERFORMANCE OF FINISHING STEERS
OFFERED MAGNESIUM-MICA IN THE
FEEDLOT RATION ¹**

K. P. Coffey² and F. K. Brazle³

Summary

Forty-eight mixed-breed steers from two sources were used in a 141-day feedlot study to compare a control ration (C) with a ration containing magnesium-mica (MM; 9 lb/ton). No diet cattle source interactions were detected. Steer gain, efficiency, and cost of gain did not differ ($P>.10$) between diets. Marbling score tended ($P<.10$) to be greater and the percentage of cattle grading USDA Choice and net carcass value were greater ($P<.05$) for steers fed MM. Feeding MM in a feedlot ration may have a substantial economic impact on feedlot cattle.

(Key Words: Magnesium-Mica, Feedlot Cattle, Marbling Score.)

Introduction

Magnesium-mica (MM) is a mined product that is used as a pellet-binding and feed-diluting agent. Previous work at KSU-SEARC has shown a tendency for increased digestibility and increased rumen fermentation products from cattle fed MM. Our objective was to measure the effect of MM on performance and carcass characteristics of feedlot cattle.

Experimental Procedures

Twenty-four mixed-breed steers previously grazing tall fescue pastures and 24 Angus Simmental crossbred steers previously grazing smooth bromegrass were allotted within cattle source into eight groups of six head each on January 20, 1994. The groups then were allotted randomly to receive one of two finishing diets consisting of 80% ground grain sorghum, 15% corn silage, and 5% protein supplement (50% CP) on a dry matter basis (Table 1). Cattle received the finishing diet for 141 days and then were slaughtered at a commercial packing plant. Carcass data were collected following a 24-hour chill.

Results and Discussion

No diet by cattle source interaction was detected in the study. Therefore, data are expressed within main effects. Steers receiving MM were somewhat heavier at the initiation of the study than were C steers (Table 2). Final weight and gain, feed cost, feed efficiency, and feed cost per cwt of gain did not differ ($P>.10$) between diets. However, total feed cost was numerically \$16.47 lower per steer and cost of gain was numerically

¹Appreciation is expressed to Micro-Lite, Inc., Chanute, KS for providing financial support; to SmithKline Beecham, West Chester, PA for providing Bovishield 4+L5, and Ultrabac/somnubac vaccinations; and to Syntex Animal Health, West Des Moines, IA for providing oxfendazole and Synovex-S.

²KSU-Southeast Agricultural Research Center, Parsons, KS.

³Southeast Kansas Area Extension Livestock Specialist, Chanute, KS.

\$3.00 less per cwt of gain for steers fed MM than for steers fed C.

Hot carcass weight, dressing percentage, fat thickness, and USDA Yield Grades did not differ ($P>.10$) between diets (Table 3). Marbling scores averaged over one third of a quality grade higher ($P<.10$) for steers fed MM. Steers in this study graded poorly. Therefore, this increase in marbling score from steers fed MM resulted in a substantial increase ($P<.05$) in the percentage of steers grading USDA Choice, which further resulted in a \$50.92 higher ($P<.05$) carcass value.

Magnesium-mica had no effect on feedlot gain, consumption, efficiency, carcass weight, or fat thickness but had a substantial effect on carcass marbling. Although this study utilized a limited number of animals (48) and marbling score is highly variable, the fact that 13 of 24 steers fed MM graded Choice (versus only 3 of 24 control steers) certainly casts doubt that the differences were due to chance. Although a biological explanation for the improved marbling score by steers fed MM is not clear, this improvement potentially represents a significant economic benefit.

Table 1. Compositions of Diets Offered to Finishing Steers

Ingredient	Control	Magnesium-Mica
Ground grain sorghum	80.0	80.0
Corn silage	15.0	15.0
Soybean meal	2.5	2.6
Wheat middlings	.81	.25
Ground limestone	.50	.50
Urea	.43	.45
TM salt	.25	.25
Cane molasses	.15	.15
Dicalcium phosphate	.13	.13
Potassium chloride	.13	.13
Vitamin A,D,E premix	.10	.10
Rumensin premix	.0125	.0125
Magnesium-mica	-	.45

Table 2. Performance by Finishing Steers Offered Magnesium-Mica (9 lb/ton) in the Feedlot Ration

Item	Diet		Cattle Source ^a	
	Control	Magnesium Mica	Purchased	SEARC
Initial wt., lb ^b	862	879	811	930
Final wt., lb ^b	1340	1354	1248	1446
Gain, lb ^c	478	475	437	516
Daily gain, lb ^c	3.4	3.4	3.1	3.7
Daily DM intake, lb ^b	29.0	28.3	26.4	30.9
Gain/feed, lb/lb	.117	.119	.118	.119
Feed cost, \$ ^c	257.25	240.78	226.20	271.83
Feed cost, \$/cwt gain	53.98	50.98	51.78	53.18

^aPurchased steers were exotic mixed-breed steers. SEARC steers were Angus Simmental and Simmental Angus crossbred steers.

^bDifferences between cattle sources were detected ($P < .05$).

^cDifferences between cattle sources were detected ($P < .10$).

Table 3. Carcass Characteristics of Finishing Steers Offered Magnesium-Mica (9 lb/ton) in the Feedlot Ration

Characteristic	Diet		Cattle Source ^a	
	Control	Magnesium Mica	Purchased	SEARC
Hot carcass wt, lb ^b	804	814	747	871
Dressing % ^c	60.0	60.0	59.8	60.3
Fat thickness, in	.30	.28	.28	.30
Longissimus eye area, in ²	13.8	13.8	13.6	14.1
Marbling score ^{def}	386	499	387	498
% USDA Choice ^{bg}	12.5	54.2	20.8	45.8
USDA Yield Grade ^b	1.4	1.6	1.2	1.8
Net carcass value, \$ ^{bg}	786.80	837.72	750.56	873.95

^aPurchased steers were exotic mixed-breed steers. SEARC steers were Angus Simmental and Simmental Angus crossbred steers.

^bDifferences between cattle sources were detected ($P < .05$).

^cCalculated using actual unshrunk live weight.

^d300 = Select⁻; 400 = Select⁰; 500 = Select⁺.

^eDifferences between diets were detected ($P < .10$).

^fDifferences between cattle sources were detected ($P < .10$).

^gDifferences between diets were detected ($P < .05$).