

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 2 *Dairy Research (1984-2014)*

Article 173

1984

Development of an early weaning program for dairy calves

J.L. Morrill

A.D. Dayton

A.J. Zmolek

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Dairy Science Commons](#)

Recommended Citation

Morrill, J.L.; Dayton, A.D.; Zmolek, A.J.; and Vitcenda, M.A. (1984) "Development of an early weaning program for dairy calves," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 2. <https://doi.org/10.4148/2378-5977.3098>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1984 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Development of an early weaning program for dairy calves

Authors

J.L. Morrill, A.D. Dayton, A.J. Zmolek, and M.A. Vitcenda

K DEVELOPMENT OF AN EARLY WEANING PROGRAM FOR DAIRY CALVES¹**S**J. L. Morrill, A. D. Dayton², A. J. Zmolek³,and M. A. Vitcenda³**U**

Summary

After preliminary experiments, four trials were conducted to develop and test a new feeding program for calves. The program involves the use of a special feed (a prestarter) to encourage calves to eat dry feed and to provide high quality nutrients during the time of initial rumen development. Using this plan, calves were weaned at 2 weeks of age with good results.

Introduction

After calves are weaned, there is a reduction in labor required, less expensive feeds are needed, and the incidence of scours usually decreases. Previous research has indicated that calves could be weaned as early as 3 wk of age with good results. In spite of this, most dairy calves in the United States are weaned at 6 to 12 wk of age. Since the main factor that determines when calves can be successfully weaned is the amount of dry feed consumed, anything that would increase dry feed consumption would help in an early weaning program. We conducted a series of experiments to evaluate the use of a special feed (a prestarter) to stimulate intake and we developed a program for the use of this prestarter.

Experimental Procedure

Preliminary observations indicated that a prestarter composed of milk solids, supplementary fat, and additives (Table 1) increased dry feed consumption if mixed with a calf starter. In Trial 1, 21 bull calves were purchased from local dairy producers and assigned to one of three treatments. Each calf was fed colostrum until 3 days of age, then milk at 8% of birth weight daily, and either prestarter, starter (Table 2), or an equal mixture of the two. Daily feed intake and weekly calf weights were recorded and the health of each calf was monitored. The calves were weaned at 2 wk of age, if consuming 1 pound of dry feed daily.

The results of Trial 1 suggested that a feeding schedule involving the use of changing ratios of prestarter and starter would be advantageous. Three subsequent trials were conducted to evaluate different feeding schedules and methods. In each, bull calves were fed milk or milk replacer and prestarter or starter according to the schedule being evaluated. Feed consumption, weight gains, fecal scores, and other observations of health were recorded.

¹Partial financial support provided by Merrick Foods, Union Center, WI.

²Department of Statistics

³Merrick Foods.

Results and Discussion

In Trial 1, calves that were fed prestarter as the only dry feed consumed the most and gained the most during the first part of the trial, those fed a mixture of prestarter and starter consumed most and gained fastest during wk 4 and 5, and calves fed only starter performed best during wk 6 (Table 3). This suggested that a program providing prestarter as the only dry feed at first, then a mixture of starter and prestarter, and finally starter alone would give good results.

During the next three trials we tested various schemes and developed the following program which gave good results when calves were weaned at 2 wk of age.

1. Ensure adequate consumption of colostrum soon after birth and during the first 3 days of life.
2. Provide adequate individual housing.
3. Feed milk or diluted colostrum at 8% of birth weight daily, using an open pail.
4. Stimulate calves to eat dry feed by putting a small amount of prestarter in the milk.
5. Provide prestarter (Table 1) ad libitum.
6. Add palatable, fibrous starter when consumption of prestarter is $\frac{1}{2}$ pound daily, first in small amounts, then increasing to appetite of calf.
7. Wean at 2 wk of age, if calf is healthy and eating dry feed well.
8. Continue to feed $\frac{1}{2}$ pound of prestarter mixed with all the starter the calf will consume to 6 wk of age.
9. Allow access to hay unless starter contains adequate roughage.

Table 1. Composition of prestarter, %

Whey, dried	46
7-60 ^a	23
Skim milk, dried	19
Sodium caseinate	12
Additives ^b	+

^a A mixture of milk solids and fat containing 7% protein and 60% animal fat.

^b Additives provided antibiotics, vitamins, and minerals.

Table 2. Composition of starter, %

Corn, ground	30
Alfalfa, ground	25
Oats, rolled	20
Soybean meal	10
Sorghum grain, rolled	7.5
Molasses, dry	5
Dicalcium phosphate	.7
Limestone, ground	.3
Salt	.25
Trace mineral salt	.25
Vitamin and mineral supplement ^a	1.00

^a Provided 1000 I.U. vitamin A, 150 I.U. vitamin D, and .4 mg selenium per pound.

Table 3. Results of Trial 1

Week	Weight gain (lb)			Feed consumed (lb)		
	Starter	50-50 mix	Prestarter	Starter	50-50 mix	Prestarter
1	3.70	4.99	6.42	.92	1.28	2.60
2	2.86	6.42	3.15	1.89	2.82	2.99
3	7.26	3.15	3.85	7.00	8.40	4.55
4	3.85	6.84	2.29	1.39	15.16	9.53
5	7.83	11.97	6.71	19.07	20.79	12.54
6	12.98	9.55	2.42	25.59	22.90	13.86