

2019

Herbicide and Application Timing Effects on Windmillgrass (*Chloris verticillata*) Control

Nicholas Mitchell

Kansas State University, nmitch2@ksu.edu

Jared Hoyle

Kansas State University, jahoye@ksu.edu

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Horticulture Commons](#), and the [Weed Science Commons](#)

Recommended Citation

Mitchell, Nicholas and Hoyle, Jared (2019) "Herbicide and Application Timing Effects on Windmillgrass (*Chloris verticillata*) Control," *Kansas Agricultural Experiment Station Research Reports*: Vol. 5: Iss. 5. <https://doi.org/10.4148/2378-5977.7768>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2019 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



TURFGRASS RESEARCH 2019



JULY 2019

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

K-State Research and Extension is an equal
opportunity provider and employer.

Herbicide and Application Timing Effects on Windmillgrass (*Chloris verticillata*) Control

Nicholas Mitchell and Jared Hoyle

Summary

Windmillgrass (*Chloris verticillata* Nutt.) populations commonly infest turfgrass systems in the midwest, which result in aesthetically unacceptable turfgrass stands. Research trials were initiated in 2017 and 2018, in Onaga and Junction City, KS, to determine windmillgrass control with various single herbicide applications at different application timings. Pylex (topramezone) resulted in more than 80% windmillgrass control 8 weeks after spring and summer application.

Rationale

Windmillgrass is a problematic perennial grassy weed commonly found in established turfgrass in the midwest. Currently, mesotrione and topramezone (4-hydroxyphenylpyruvate dioxygenase [HPPD] inhibitor) are the only available labeled post-emergent active ingredients for windmillgrass control in turfgrass systems.

Objective

The objective of this study was to determine the effect of a single post-emergent herbicide application and optimal application timing for windmillgrass control.

Study Description

Research trials were initiated in 2017 in Onaga and Junction City, KS, to determine windmillgrass herbicide efficacy as well as application timing to provide windmillgrass control. Windmillgrass populations consisted of 65–75% cover in a low maintenance tall fescue (*Schedonorus arundinaceus*) system maintained at 2.5 inches. No supplemental irrigation was applied to the research areas throughout the duration of the trials. Treatments were arranged in a 3 × 10 factorial, randomized complete

View all turfgrass research reports online at: <http://newprairiepress.org/kaesrr>



block design with 4 replications. Factors were the combination of 3 application timings (spring, summer, and fall) and 10 herbicides. Summer treatments were applied on July 14, 2017, in Junction City, KS, and on July 19, 2017, in Onaga, KS. The fall treatments were applied on September 7, 2017, in Junction City, KS, and on September 15, 2017, in Onaga, KS. The spring treatments were applied on May 24, 2018, in Junction City, KS, and May 25, 2018, in Onaga, KS. Herbicide treatments consisted of Tenacity (mesotrione) at 8 fl oz/a, Acclaim (fenoxaprop) at 39 fl oz/a, Pylex (topramezone) at 2 fl oz/a, Drive XLR8 (quinclorac) at 64 fl oz/a, Dismiss (sulfentrazone) at 8 fl oz/a, Revolver (foramsulfuron) at 32.5 fl oz/a, Celsius WG (thiencarbazone + iodosulfuron + dicamba) at 4.9 oz/a, Katana (flazasulfuron) at 3 oz/a, Monument 75WG (trifloxysulfuron) at 0.53 oz/a, and a nontreated control. A surfactant was added to each herbicide treatment according to the manufacturer's directions. Herbicide treatments were applied using a CO₂ pressurized backpack boom sprayer calibrated to deliver 43.56 GPA at 37 psi. Data collected for research trials consisted of visual percent windmillgrass cover (0–100%). Evaluations were conducted weekly for 4 weeks and then biweekly to 8 weeks after each application. Analysis of variance (ANOVA) was performed in SAS 9.4 (SAS Institute Inc., Cary, NC) and means were separated according to Fisher's protected least significant difference (LSD) level at 0.05.

Results

Pylex (topramezone) applied at 2 fl oz/a in the spring and summer resulted in 83% and 81% windmillgrass control at 8 weeks after treatment, respectively. Spring and summer applications of Pylex at Rolling Meadows resulted in much greater control compared to the fall application. Drive (quinclorac), Dismiss (sulfentrazone), Revolver (foramsulfuron), Celsius (thiencarbazone + iodosulfuron + dicamba), Katana (flazasulfuron), and Monument (trifloxysulfuron) resulted in unacceptable windmillgrass control ($\leq 13\%$) with spring and summer applications at both research locations. No herbicides tested in this study applied in the spring, summer, or fall with a single application resulted in acceptable windmillgrass control.

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service





Table 1. Windmillgrass control by various herbicide treatments following spring, summer, and fall applications in Junction City, KS

Herbicide Treatment	Application timing ^a		
	Spring	Summer	Fall
	% Control ^d		
Tenacity	15 bc ^b , B ^c	9 bc, B	43 a, A
Acclaim	31 b, A	30 b, A	43 a, A
Pylex	83 a, A	81 a, A	54 a, B
Drive	9 c, B	0 c, B	47 a, A
Dismiss	0 c, B	0 c, B	35 ab, A
Revolver	4 c, B	0 c, B	43 a, A
Celsius	5 c, B	0 c, B	34 abc, A
Katana	4 c, A	0 c, A	15 bc, A
Monument	13 bc, A	0 c, A	11 c, A

^aRatings were done 8 weeks after each application: July 19, 2018 (spring), September 7, 2017 (summer), and November 2, 2017 (fall).

^bTreatment means followed by a common lowercase letter are not significantly different within application timing according to Fisher's protected LSD ($\alpha = 0.05$).

^cTreatment means followed by a common capital letter are not significantly different within active ingredient according to Fisher's protected LSD ($\alpha = 0.05$).

^dNon-treated control contained 50, 65, and 45% windmillgrass cover at 8 weeks after treatment for spring, summer, and fall application timings, respectively. % Control = $[(A-B)/A] \times 100$, A = % windmillgrass cover at applications and B = % windmillgrass cover 8 weeks after application.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

