

# Kansas Agricultural Experiment Station Research Reports

---

Volume 10  
Issue 5 *Western Kansas Agricultural Research*

Article 17

---

2024

## Single and Split Applications for Herbicide Efficacy in Corn

Patrick W. Geier

Kansas State University, [pgeier@k-state.edu](mailto:pgeier@k-state.edu)

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Agronomy and Crop Sciences Commons](#), and the [Weed Science Commons](#)

---

### Recommended Citation

Geier, Patrick W. (2024) "Single and Split Applications for Herbicide Efficacy in Corn," *Kansas Agricultural Experiment Station Research Reports*: Vol. 10: Iss. 5. <https://doi.org/10.4148/2378-5977.8614>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2024 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



## Single and Split Applications for Herbicide Efficacy in Corn

*P. W. Geier*

### Summary

This study investigated the use of residual herbicides as single or sequential applications for efficacy in corn. All preemergence (PRE) herbicides provided good control of Palmer amaranth and green foxtail early in the season. Storen plus atrazine PRE and all PRE followed by postemergence (POST) treatments controlled Palmer amaranth best later in the season. Lumax EZ plus atrazine and Zidua SC plus Callisto PRE were the only treatments to provide less than 90% green foxtail control later in the year. Late-season Johnsongrass control was best with sequential applications of Acuron or Storen (90 to 96%).

### Introduction

The use of residual herbicides early during crop establishment is an important component for preserving yield later in the season. If part of that residual herbicide can be applied later with a planned POST application, the weed-free period can be extended, allowing the crop to become established and better compete for resources. The objective of this trial was to compare single and sequential applications of residual herbicides for efficacy and crop yield in corn.

### Experimental Procedures

An experiment compared the use of residual herbicides as either a single or as sequential applications for weed control in corn. Herbicides were applied using a tractor-mounted, compressed-CO<sub>2</sub> sprayer delivering 19.4 gpa at 30 psi and 4.1 mph. Application, environmental, and plant information is shown in Table 1. Plots were 10 by 35 ft and arranged in a randomized complete block design replicated four times. Soil was Beeler silt loam having 2.4% organic matter, pH of 7.5, and CEC of 17.8. Visual weed control was determined on May 29 and July 10, 2023. These dates were 24 days after the preemergence applications (24 DAA) and 42 days after the POST applications (42 DAB), respectively. Corn yields were determined on October 9, 2023, by mechanically harvesting the center two rows of each plot and adjusting grain moistures to 15.5%.

### Results and Discussion

All herbicides provided excellent control of kochia and Russian thistle regardless of evaluation date (data not shown). Likewise, all PRE herbicides controlled Palmer amaranth and green foxtail 98% or more at 24 DAA (Table 2). By 42 DAB, Lumax EZ, Acuron, or Storen applied as sequential treatments controlled Palmer amaranth 90 to

96%, whereas the PRE treatments of Storen or Acuron provided 85% to 89% control. Only Lumax EZ plus atrazine PRE and Zidua SC plus Callisto PRE provided less than 90% green foxtail control at 42 DAB. Johnsongrass control ranged from 70% to 88% at 24 DAA. By 42 DAB, sequential applications of Acuron or Storen were the only treatments to control johnsongrass by as much as 90%. Of the PRE-only treatments, Storen plus atrazine and Zidua SC plus atrazine were more effective on Johnsongrass late in the season than the other treatments. All herbicide-treated corn yielded more grain than nontreated corn. However, corn receiving sequential applications of Lumax EZ, Acuron, or Storen produced the highest yields.

## Acknowledgments

Funding for this research was provided by Syngenta AG.

**Table 1. Application, environmental, and plant information in the single and split application study in corn**

| Application timing                | Preemergence   | Postemergence |
|-----------------------------------|----------------|---------------|
| Application date                  | May 5, 2023    | May 29, 2023  |
| Air temperature (F)               | 62             | 67            |
| Relative humidity (%)             | 86             | 75            |
| Soil temperature (F)              | 60             | 62            |
| Wind speed (mph)                  | 4 to 7         | 3 to 6        |
| Wind direction                    | East-southeast | South         |
| Soil moisture                     | Good           | Good          |
| Corn                              |                |               |
| Height (inches)                   | ---            | 5 to 7        |
| Leaves (no.)                      | 0              | 2 to 3        |
| Russian-thistle                   |                |               |
| Height (inches)                   | ---            | 1 to 4        |
| Density (plants/ft <sup>2</sup> ) | 0              | 0.1           |
| Palmer amaranth                   |                |               |
| Height (inches)                   | ---            | 0.25 to 1     |
| Density (plants/ft <sup>2</sup> ) | 0              | 0.2           |
| Johnsongrass                      |                |               |
| Height (inches)                   | ---            | 1 to 3        |
| Density (plants/ft <sup>2</sup> ) | 0              | 0.5           |
| Kochia                            |                |               |
| Height (inches)                   | ---            | 0.5 to 2      |
| Density (plants/ft <sup>2</sup> ) | 0              | 0.1           |
| Green foxtail                     |                |               |
| Height (inches)                   | ---            | 0.5 to 1      |
| Density (plants/ft <sup>2</sup> ) | 0              | 0.1           |

Table 2. Weed control and grain yield in the single and split application study in corn

| Treatment            | Rate   | Timing <sup>1</sup> | Palmer amaranth      |                     | Green foxtail |        | Johnsongrass |        | Corn yield |
|----------------------|--------|---------------------|----------------------|---------------------|---------------|--------|--------------|--------|------------|
|                      |        |                     | 24 DAA <sup>2</sup>  | 42 DAB <sup>2</sup> | 24 DAA        | 42 DAB | 24 DAA       | 42 DAB |            |
|                      | qt/a   |                     | ----- % Visual ----- |                     |               |        |              |        | bu/a       |
| Untreated            | ---    | ---                 | ---                  | ---                 | ---           | ---    | ---          | ---    | 23.1       |
| Lumax EZ             | 2.7    | PRE                 | 100                  | 73                  | 100           | 83     | 75           | 58     | 55.4       |
| Atrazine             | 0.5    | PRE                 |                      |                     |               |        |              |        |            |
| Acuron               | 3.0    | PRE                 | 100                  | 85                  | 100           | 98     | 78           | 60     | 70.8       |
| Atrazine             | 0.5    | PRE                 |                      |                     |               |        |              |        |            |
| Storen               | 2.1    | PRE                 | 100                  | 89                  | 100           | 90     | 85           | 80     | 85.6       |
| Atrazine             | 1.25   | PRE                 |                      |                     |               |        |              |        |            |
| Lumax EZ             | 1.35   | PRE                 | 100                  | 90                  | 100           | 100    | 70           | 88     | 113.8      |
| Atrazine             | 0.25   | PRE                 |                      |                     |               |        |              |        |            |
| Lumax EZ             | 1.35   | POST                |                      |                     |               |        |              |        |            |
| Atrazine             | 0.25   | POST                |                      |                     |               |        |              |        |            |
| Glyphosate           | 25 oz  | POST                |                      |                     |               |        |              |        |            |
| Ammonium sulfate     | 2.5 %  | POST                |                      |                     |               |        |              |        |            |
| Acuron               | 1.5    | PRE                 | 100                  | 93                  | 98            | 100    | 75           | 90     | 122.5      |
| Atrazine             | 0.25   | PRE                 |                      |                     |               |        |              |        |            |
| Acuron               | 1.5    | POST                |                      |                     |               |        |              |        |            |
| Atrazine             | 0.25   | POST                |                      |                     |               |        |              |        |            |
| Glyphosate           | 25 oz  | POST                |                      |                     |               |        |              |        |            |
| Ammonium sulfate     | 2.5 %  | POST                |                      |                     |               |        |              |        |            |
| Storen               | 1.05   | PRE                 | 100                  | 96                  | 100           | 100    | 80           | 96     | 138.2      |
| Atrazine             | 0.63   | PRE                 |                      |                     |               |        |              |        |            |
| Storen               | 1.05   | POST                |                      |                     |               |        |              |        |            |
| Atrazine             | 0.63   | POST                |                      |                     |               |        |              |        |            |
| Glyphosate           | 25 oz  | POST                |                      |                     |               |        |              |        |            |
| Ammonium sulfate     | 2.5 %  | POST                |                      |                     |               |        |              |        |            |
| Zidua SC             | 3.5 oz | PRE                 | 98                   | 75                  | 98            | 83     | 88           | 73     | 105.6      |
| Atrazine             | 1.25   | PRE                 |                      |                     |               |        |              |        |            |
| Degree Xtra          | 3.0    | PRE                 | 100                  | 80                  | 100           | 90     | 78           | 55     | 75.2       |
| Callisto             | 5.0 oz | PRE                 |                      |                     |               |        |              |        |            |
| Bicep Lite II Magnum | 1.67   | PRE                 | 100                  | 65                  | 100           | 95     | 78           | 55     | 53.3       |
| Callisto             | 5.0 oz | PRE                 |                      |                     |               |        |              |        |            |
| LSD (0.05)           |        |                     | NS                   | 6                   | NS            | 12     | 10           | 9      | 27.9       |

<sup>1</sup> PRE = preemergence. POST = postemergence.<sup>2</sup> DAA = days after the preemergence application. DAB is days after the postemergence application.