

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 10 *Swine Day (1968-2014)*

Article 90

1972

Effects of flaked and roasted sorghum grain on finishing swine performance

G L. Allee

Robert H. Hines

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Allee, G L. and Hines, Robert H. (1972) "Effects of flaked and roasted sorghum grain on finishing swine performance," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 10. <https://doi.org/10.4148/2378-5977.3510>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1972 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



K**S****U**

Effects of Flaked and Roasted Sorghum Grain on Finishing Swine Performance

R. H. Hines and Gary L. Allee

Summary

The value of feeding flaked or roasted sorghum grain to finishing swine (95 to 210 lbs.) was investigated using 54 barrows and gilts. Rate and efficiency of gain for each treatment group were similar. Data from this station indicate that processing methods like flaking or roasting sorghum grain do not improve the performance of finishing swine enough to offset the increased cost of processing.

Procedures

Fifty-four barrows and gilts (Hampshire, Yorkshire, and Duroc) averaging 95 lbs. were allotted to three treatments: a) ground sorghum grain (basal ration); b) flaked sorghum grain; c) roasted sorghum grain.

The flaked grain was prepared by steam heating followed by flat rolling and cooling. The roasted grain was prepared by use of a commercial roast-a-tron machine. The complete ration consisted of 1600 lbs. of processed grain plus 400 lbs. of protein supplement (table 1). Each complete ration contained approximately 16% crude protein, 0.63% phosphorous, and 0.78% calcium. The flaked and ground rations contained approximately 14% moisture; the roasted grain, 12%.

Pigs were housed in an open-front barn, with slatted pens, 6' x 15'. Each pen of 9 pigs had access to a two-hole self feeder and an automatic watering cup.

The feeding trial was for 70 days during the summer months.

Results and Discussion

Performance data are shown in table 5. Finishing pigs (95 to 210 lbs.) gained at a similar rate whether they were fed ground, flaked, or roasted sorghum grain. Feed:gain ratios for the three processes were also similar.

The 1972 results confirm the 1971 results with processed milo for finishing swine (Report of Progress 181, Swine Industry Day, 1971).

Table 5. Performance Data of Finishing Swine Fed Ground, Flaked, or Roasted Sorghum Grain

Ration:	<u>Ground</u>	<u>Flaked</u>	<u>Roasted</u>
<u>Daily gain, lbs.</u>			
Rep. 1	1.52	1.45	1.45
Rep. 2	<u>1.38</u>	<u>1.37</u>	<u>1.50</u>
Average	1.45	1.41	1.47
<u>Feed intake, lbs./day</u>			
Rep. 1	4.92	4.84	4.82
Rep. 2	<u>4.82</u>	<u>4.62</u>	<u>4.94</u>
Average	4.87	4.73	4.88
<u>Feed/gain</u>			
Rep. 1	3.24	3.34	3.32
Rep. 2	<u>3.48</u>	<u>3.36</u>	<u>3.29</u>
Average	3.36	3.35	3.31
