

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1161

1982

Feeding Rumensin to yearling heifers on late-summer grass

M. Jensen

R. Held

E.F. Smith

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Jensen, M.; Held, R.; Smith, E.F.; and Riley, Jack G. (1982) "Feeding Rumensin to yearling heifers on late-summer grass," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2564>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1982 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Feeding Rumensin to yearling heifers on late-summer grass

Authors

M. Jensen, R. Held, E.F. Smith, and Jack G. Riley

K**S****U**

Feeding Rumensin^R to Yearling Heifers on Late-summer Grass

Mike Jensen, Rosalie Held, Jack Riley and E.F. Smith

Summary

Rumensin was self-fed in a loose salt mixture to yearling heifers on late-summer bluestem pasture (August to November). They ate 0.036 lbs. of salt mixture (124 mg of Rumensin) daily, but performance did not improve.

Introduction

Rumensin has been shown to improve the gain and efficiency of grazing cattle. Its intake has been regulated in a variety of ways. We tested regulating Rumensin intake in a loose salt mixture.

Experimental Procedure

Thirty-two Angus, Hereford, and crossbred heifers were grazed in four groups from August 4 to November 4, 1981. Two groups received a salt/Rumensin mixture and the other two were fed only salt. The cattle were allotted into four pastures, at an average stocking rate of 8 acres per head. The pastures had not been grazed since spring. The heifers were rotated between pastures every 2 weeks. The mixtures were fed free-choice in commercial mineral feeders. The amount fed was weighed and adjusted weekly according to intake. The cattle were weighed at the first of each month, after being penned overnight without feed or water.

Results and Discussion

Average daily gains of 0.72 pounds for Rumensin-group steers were statistically similar to the 0.67 pounds for the controls. The average daily Rumensin intake ranged from 18.4 to 200.5 mg. Adding Rumensin to the salt decreased salt intake by 22.8%. The wide variation of salt intake on the dry, mature grass prevented the close regulation of Rumensin intake. Also, the maturity of the late-summer grass contributed to decreased gain and could have affected the response.

^RRumensin, product of Elanco Products Co., Indianapolis, IN 46706.