

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 1062

1985

Sodium bicarbonate and sodium bentonite supplements for cattle fed corn or sorghum silages

D. Axe

K. Bolsen

K. Jacques

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Axe, D.; Bolsen, K.; Jacques, K.; and Harmon, D. (1985) "Sodium bicarbonate and sodium bentonite supplements for cattle fed corn or sorghum silages," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.2465>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 1985 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



Sodium bicarbonate and sodium bentonite supplements for cattle fed corn or sorghum silages

Authors

D. Axe, K. Bolsen, K. Jacques, and D. Harmon

K**S****U**

Sodium Bicarbonate and Sodium Bentonite Supplements for Cattle Fed Corn or Sorghum Silages^{1,2}

Dirk Axe, Keith Bolsen, Kate Jacques,
and Dave Harmon

Summary

Supplementing forage sorghum silage rations with sodium bicarbonate (NaHCO_3) improved performance of growing cattle over the control supplement. However, neither NaHCO_3 nor sodium bentonite supplementation to corn silage rations improved cattle performance.

Introduction

Beef cattle research with sodium bicarbonate (NaHCO_3) has involved mainly high concentrate feeding programs with little emphasis on silage-based rations for growing/backgrounding cattle. In two previous trials at Manhattan, addition of NaHCO_3 to high silage rations has improved rate and efficiency of gain (Reports of Progress 427 and 448). Sodium bentonite (colloid clay), an inert material, is not new to the cattle feeding industry, but results have been inconsistent. These trials further evaluated NaHCO_3 and sodium bentonite supplements for growing cattle fed forage sorghum and corn silage rations.

Experimental Procedures

Trial 1. Forage sorghum silage (Pioneer 947) was fed to 36 individually housed calves for 84 days, beginning November 17, 1983. Eighteen calves per treatment received supplements containing either no additive (control) or NaHCO_3 fed at 1.0% of the ration dry matter (DM) intake (approximately 43 grams per calf daily). The silages were full-fed and all calves received 1.8 lb of supplement daily (DM basis). The rations were formulated to provide 12.5% crude protein, 150 mg of Rumensin[®] per calf daily, and NRC recommended amounts of calcium, phosphorus, and vitamins A, D, and E.

Trial 2. Drought-stressed and irrigated whole-plant corn silages were fed to light weight yearling steers and heifers for 84 days, beginning February 9, 1984. Eight pens of four cattle were fed supplements with: 1) no additive (control), 2) NaHCO_3 , and 3) sodium bentonite. NaHCO_3 was fed at 1% of the ration DM intake (about 66 grams per animal daily), and sodium bentonite, at 2% of the ration DM intake (about 132 g per animal daily). Silages were full-fed and all cattle received

¹The sodium bicarbonate and partial financial assistance. were provided by Church and Dwight Co., Inc., Piscataway, NJ.

²The sodium bentonite was supplied by American Colloid Co., Skokie, IL.

1.8 lb of supplement daily (DM basis). Rations were formulated to provide 12% crude protein, 200 mg of Rumensin® per animal daily, and NRC recommended amounts of calcium, phosphorus, and vitamins A, D, and E.

Supplements in both trials were top-dressed and partially mixed with the silages in the bunk. All calves were weighed individually on two consecutive days at the start and at the end of the trials. Intermediate weights were taken at 28 and 56 days.

Results

Shown in Table 25.1 are performance results of the cattle in trial 1. The 0 to 84 days results show that NaHCO_3 improved rate of gain (8.8%), feed intake (2.0%), and efficiency of gain (7.5%) over the control supplement. However, the differences were not statistically significant. During days 0 to 28, the NaHCO_3 supplement gave an advantage in performance and during days 29 to 56, a period of extremely cold weather, NaHCO_3 produced a significant response in gain and feed/gain over the control supplement. A similar response was observed a year earlier (Report of Progress 448) under comparable cold weather conditions. There was some compensating performance for cattle fed the control supplement in the final 57 to 84 days.

Shown in Table 25.2 are performance results for trial 2. In general, neither NaHCO_3 nor sodium bentonite supplementation gave a performance advantage over the control supplement. Cattle fed sodium bentonite gained slower ($P < .05$) and were 4.7% less efficient than those fed the control.

Table 25.1. Performance by Cattle Fed Control and NaHCO_3 Supplements in Trial 1

Item	Control	NaHCO_3
No. of Calves	18	18
Initial Wt., lb	467	467
Final Wt., lb	553	560
	0 to 28 days	
Avg. Daily Gain, lb	1.37	1.45
Avg. Daily Feed, lb ¹	9.85	9.97
Feed/lb of Gain, lb ¹	7.6	7.1
	29 to 56 days	
Avg. Daily Gain, lb	.52 ^b	.83 ^a
Avg. Daily Feed, lb ¹	11.86	12.25
Feed/lb of Gain, lb ¹	31.8 ^b	18.6 ^a
	57 to 84 days	
Avg. Daily Gain, lb	1.16	1.05
Avg. Daily Feed, lb ¹	12.87	12.85
Feed/lb of Gain, lb ¹	12.3	15.8
	0 to 84 days	
Avg. Daily Gain, lb	1.02	1.11
Avg. Daily Feed, lb ¹	11.48	11.71
Feed/lb of Gain, lb ¹	11.8	10.9

^{a b}Values in the same row with different superscripts differ ($P < .05$).

¹ 100% dry matter basis.

Table 25.2. Performance by Cattle Fed Control, NaHCO₃, and Sodium Bentonite Supplements in Trial 2

Item	Control	NaHCO ₃	Sodium Bentonite
No. of Cattle	32	32	32
Initial Wt., lb	482	474	476
Final Wt., lb	673	663	659
0 to 28 days			
Avg. Daily Gain, lb	1.69	1.72	1.77
Avg. Daily Feed, lb ¹	12.66 ^a	13.00 ^{ab}	13.21 ^b
Feed/lb of Gain, lb ¹	7.6	7.6	7.5
29 to 56 days			
Avg. Daily Gain, lb	2.37	2.28	2.17
Avg. Daily Feed, lb ¹	14.75	14.98	14.96
Feed/lb of Gain, lb ¹	6.3	6.6	7.1
57 to 84 days			
Avg. Daily Gain, lb	2.82	2.73	2.59
Avg. Daily Feed, lb ¹	15.57	15.41	15.93
Feed/lb of Gain, lb ¹	5.5 ^a	5.6 ^a	6.1 ^b
0 to 84 days			
Avg. Daily Gain, lb	2.28 ^a	2.25 ^{ab}	2.18 ^b
Avg. Daily Feed, lb ¹	14.51	14.34	14.61
Feed/lb of Gain, lb ¹	6.4	6.4	6.7

^{ab}Values in the same row with different superscripts differ (P<.05).

¹100% dry matter basis.