

Kansas Agricultural Experiment Station Research Reports

Volume 0
Issue 1 *Cattleman's Day (1993-2014)*

Article 134

2007

The effects of ractopamine-HCl (Optaflexx) on finishing feedlot heifers

M.J. Quinn

E.R. Loe

M.E. Corrigan

See next page for additional authors

Follow this and additional works at: <https://newprairiepress.org/kaesrr>



Part of the [Other Animal Sciences Commons](#)

Recommended Citation

Quinn, M.J.; Loe, E.R.; Corrigan, M.E.; Drouillard, James S.; and Depenbusch, Brandon E. (2007) "The effects of ractopamine-HCl (Optaflexx) on finishing feedlot heifers," *Kansas Agricultural Experiment Station Research Reports*: Vol. 0: Iss. 1. <https://doi.org/10.4148/2378-5977.1537>

This report is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Kansas Agricultural Experiment Station Research Reports by an authorized administrator of New Prairie Press. Copyright 2007 the Author(s). Contents of this publication may be freely reproduced for educational purposes. All other rights reserved. Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. K-State Research and Extension is an equal opportunity provider and employer.



The effects of ractopamine-HCl (Optaflexx) on finishing feedlot heifers

Authors

M.J. Quinn, E.R. Loe, M.E. Corrigan, James S. Drouillard, and Brandon E. Depenbusch

THE EFFECTS OF RACTOPAMINE-HCL (OPTAFLEXX¹) ON FINISHING FEEDLOT HEIFERS

M. J. Quinn, J. S. Drouillard, E. R. Loe, B. E. Depenbusch, and M. E. Corrigan

Introduction

Whether occurring naturally or synthetically, beta-agonists are classified as phenethanolamines based on their chemical structure. Other specificities of structure determine the exact behavior of the compound in animals. Beta-agonists bind to specific receptors on the cell membranes of skeletal muscle, adipose, and other tissues. Upon binding, these agents alter metabolic pathways, ultimately causing a repartitioning of nutrients to increase muscle and decrease fat accretion. In cattle, beta-agonists such as cimaterol and clenbuterol improve feed efficiency, average daily gain, and longissimus muscle area. The proposed effects of ractopamine HCl are similar to those of other beta-agonists. The purpose of this study was to examine the effects of ractopamine-HCl on live performance, carcass characteristics, and meat quality of finishing beef heifers.

Experimental Procedures

Three hundred and two crossbred heifers (890±3 lbs) were purchased from salebarns and transported to Kansas State University Beef Cattle Research Center in Manhattan. All cattle were offered *ad libitum* access to hay and water prior to processing. Body weight was determined and the animals were given an internal/external parasiticide. Heifers were implanted with Revalor²-H (140 mg TBA, 14

mg estradiol) and gradually adapted to a diet containing 94% concentrate and 6% alfalfa hay. At the end of the adaptation phase, the heifers were blocked by initial weight and allotted to dirt-surfaced feeding pens containing 12 to 13 animals each. Treatments were with Optaflexx and a control treatment. Cattle receiving the Optaflexx treatment were fed 200 mg/day of ractopamine-HCl, beginning 28 days prior to slaughter. Pens of cattle were weighed using a platform scale on day 0, at approximately 28-day intervals thereafter, prior to the Optaflexx period, and immediately before being transported to a commercial abattoir for slaughter.

All cattle were allowed *ad libitum* access to a common finishing diet. The entire daily ration was delivered at approximately 1:00 p.m. At the end of the experiment, heifers were shipped to a commercial abattoir in Emporia, Kansas. Slaughter data, including hot carcass weight, incidence and severity of liver abscess, and dress yield, were obtained on the day of slaughter. Following a 24-hour chill, carcasses were evaluated for subcutaneous fat thickness; kidney, pelvic, and heart fat; longissimus muscle area; marbling score; and USDA yield and quality grades.

Following fabrication, loins were obtained from three animals randomly selected from each pen. The loins were aged for 14 days in

¹Optaflexx is a registered trademark of Elanco Animal Health, Indianapolis, IN.

²Revalor is a registered trademark of Intervet, Inc.

cryovac packages at 32±2 F. After aging, loin steaks were removed and vacuum packaged for sampling. Loin steak samples were analyzed for fatty acid composition, purge loss measurements from a seven-day display period, and Warner-Bratzler shear force. Loin steak samples were cooked using to an internal temperature of 160° F. Samples were then refrigerated at 38 F for 24 hours. Following the 24-hour refrigeration period, six to eight half-inch cores were removed parallel to the fiber orientation of the steaks for shear force analysis.

Results and Discussion

Over the entire 75-day feeding period, dry matter intake, average daily gain, and efficiency of gain were not different between heifers that received 0 or 200 mg/day of ractopamine-HCl ($P>0.17$). However, Optaflexx-fed cattle had improved feed efficiency for the 28-day Optaflexx period ($P = 0.056$). Similar measurements ($P>0.19$) were obtained for the two treatments for dressing percent; hot carcass weight; marbling score; fat thickness; ribeye area; kidney, pelvic and heart fat; and USDA yield and quality grades. There was also no difference in Warner-Bratzler shear force between the two treatments ($P>0.40$). Purge loss during the seven day commercial display period and loss during cooking also

were not different between treatments ($P>0.50$).

Table 1. Ingredients and Nutrient Composition of Finishing Diets

Item, % dry basis	Diet
Steam-flaked corn	79.55
Ground alfalfa hay	6.00
Corn-steep liquor	6.23
Vitamin/mineral supplement	6.01
Feed additive premix ^a	2.21 ^a
Nutrient	
Dry matter	80.0
Crude protein	14.00
Fat	3.70
Calcium	0.75
Phosphorus	0.39

^aFormulated to provide 300 mg monensin, 90 mg Tylosin, 0.5 mg MGA, and 0 or 200 mg ractopamine-HCl daily in a ground corn carrier.

Implications

Optaflexx added to the diets of finishing beef heifer improved gain efficiency during the 28-day feeding period with no effect on carcass quality or meat characteristics.

Table 2. Live Performance of Heifers Fed 0 or 200 mg Ractopamine-HCl for 28 Days Prior to Finishing

Item	Control	Optaflexx	SEM	P-Value
Number of heifers	150	152	-	-
Days fed Optaflexx	0	28	-	-
Initial weight, lb	1055	1054	6	0.91
Daily dry matter intake, lb	22.2	21.7	0.25	0.19
Average daily gain, lb	3.72	3.95	0.12	0.17
Feed:gain	5.98	5.46	0.005	0.06
Final weight ^a , lb	1159	1164	7	0.59

^aFinal body weight based on carcass adjusted weight (carcass weight divided by 63.5% dress).

Table 3. Carcass Characteristics of Heifers Fed Either 0 or 200 mg/day of Ractopamine HCl

Characteristic	Control	Optaflexx	SEM ²	P-Value
Carcass weight, lb	736	739	5	0.59
Ribeye area, square inches	13.38	13.67	0.2	0.31
Fat thickness (12 th rib), inches	0.31	0.30	0.01	0.47
Kidney, pelvic, heart fat, %	2.24	2.23	0.03	0.85
Marbling ¹	380	373	7	0.48
Average yield grade	1.99	2.01	0.07	0.85
USDA Choice, %	36.2	40.7	5.1	0.55
USDA Select, %	57.8	56.0	5.4	0.81
USDA Standard, %	6.0	3.4	2.0	0.37
USDA YG1, %	26.4	21.2	4.4	0.41
USDA YG2, %	51.6	58.3	4.2	0.28
USDA YG3, %	17.2	19.2	3.1	0.65
USDA YG4, %	4.1	1.3	1.5	0.19

¹Traces=200-299; Slight=300-399; Small=400-499.

²SEM=Standard Error of the Mean.

Table 4. Meat Quality Characteristics of Control and Optaflexx Fed Heifers

Item	Control	Optaflexx	SEM	P-Value
Warner-Bratzler shear force, lbs	10.1	10.4	1.1	0.41
Purge loss from retail display, %	3.2	3.3	0.7	0.58
Purge loss from cooking, %	25	25	14	0.51
L*	42.4	41.5	5	0.11
a*	31.0	30.0	5	0.40
b*	24.1	23.7	2	0.17